

Summit ... responsible solutions.

Resealable Label **OPEN HERE** for Directions & Precautions

Biological Caterpillar & Webworm Control

FOR ORGANIC GARDENING



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **CAUTION:**

Causes moderate eye irritation. Avoid contact with eyes or clothing Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor, or going for treatment

Hot Line Number: 1-800-255-3924.



ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

Summit Biological Caterpillar & Webworm Control is an easy to mix liquid concentrate.

- Kills worms and caterpillar stage insects, but has no effect on birds, earthworms, or beneficial insects, such as
- honeybees and ladybugs, when used as directed.
- · Worms and caterpillars eat treated foliage, then immediately stop feeding and damaging plants. · Acceptable for use on edible plants up to the day of harvest.

DIRECTIONS FOR USE

It is a violation of Federal I aw to use this product in a manner inconsistent with its labeling

Apply Summit Biological Caterpillar & Webworm Control when worms or caterpillars are first noticed, then repeat at five (5) to seven (7) day intervals while they are active. Apply more frequently to control heavy infestations Apply thoroughly to top and bottom of foliage. Reapply after heavy rains.

Summit Biological Cateroillar & Webworm Control must be eaten by worms or cateroillars to be effective After ingesting the insecticide, they immediately stop feeding, though they may otherwise appear to be unaffected for several days. Best results are obtained by treatments when worms are small; they must be actively feeding on treated, exposed foliage. You may apply Summit Biological Caterpillar & Webworm Control up to and on the day of harvest

How to Apply: Always shake or stir Summit Biological Caterpillar & Webworm Control for Home and Gardens thoroughly before use. Partially fill sprayer with water before adding the application rate amount of product, then mix in product thoroughly and add remaining amount of water

Spray leaf surfaces thoroughly, top and bottom, for complete control, Agitate regularly while spraying, Use all of spray mixture within 24 hours

Fruits and Vegetables: Mix 1.5 fluid ounces per 3 gallons water (1 tablespoon per gallon) for a hand sprayer and apply to all plant foliage to cover an area of 1000 square feet. Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control of the following insects:

Cabbage Looper on broccoli, cauliflower. collards, kale, mustard greens, turnip greens, cabbage, celery, lettuce, melons, and tomatoes,

Imported Cabbage Worm on broccoli. cabbage cauliflower collards kale mustard greens, and turnip greens.

Tomato Hornworm on tomatoes

MIXING DIRECTIONS

Make sure the spray equipment to be used is clean, dry and odor free.

The attached sprayer is calibrated to deliver a 1 to 100 dilution.

Hose-end convenience!
Automatically mixes with water! No measuring or separate applicator!



. Attach garden







Shade Trees & Ornamentals: Mix 2 fluid ounces per 3 gallons of water (4 teaspoons per gallon) for a hand sprayer or proportional amounts for a hose-end sprayer following manufacturer's directions.

Apply at first sign of infestation and repeat at weekly intervals when needed to maintain control. Apply to thoroughly cover all foliage surfaces of shade trees and ornamentals for control of the following leaf-feeding worms

Bagworm Spring Cankerworm Fall Cankerworm Gypsy Moth* Tent Caterpillar Elm Spanworm

* Apply when buds swell, but before leaves are fully open. If eggs hatch over a long period of time or if reinfestation occurs, spray about 14 days after first application

Fall Webworm

STORAGE AND DISPOSAL

Pesticide Storage: Store in a cool place. Activity may be impaired by storage at temperatures above 90°F,

Pesticide Disposal and Container Handling:

Non-refillable container, Do not reuse or refill this container,

If empty:

Place in trash or offer for recycling if available,

Call your local solid waste agency or 1-800- CLEANUP for disposal instructions, Never place unused product down any indoor or outdoor drain,

WARRANTY

Summit Chemical Company warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions

NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OF MERCHANTABILITY IS MADE.



235 South Kresson Street Baltimore, MD 21224-2616 800-227-8664

www.SummitChemical.com

SUMMIT RESPONSIBLE SOLUTIONS® IS A REGISTERED TRADEMARK OF SUMMIT CHEMICAL CO., BALTIMORE, MARYLAND,